

sub  
A1

[illegible]

1. A method for identifying one or more low abundance sequences differing by one or more single-base changes, insertions, or deletions, from a high abundance sequence in a plurality of target nucleotide sequences comprising:
- 5 providing a sample potentially containing one or more low abundance target nucleotide sequences with at least one sequence difference each from the high abundance target sequences;
- providing a primary oligonucleotide primer set characterized by (a)
- 10 a first oligonucleotide primer containing a target-specific portion, and (b) a second oligonucleotide primer containing a target-specific portion, wherein the primary oligonucleotide primers are suitable for hybridization on complementary strands of a corresponding high and low abundance target nucleotide sequences to permit formation of a polymerase chain reaction product, but have a mismatch which
- 15 interferes with formation of such a polymerase chain reaction product when hybridized to any other nucleotide sequence present in the sample;
- providing a polymerase;
- blending the sample, the primary oligonucleotide primers, and the polymerase to form a primary polymerase chain reaction mixture;
- 20 subjecting the primary polymerase chain reaction mixture to two or more polymerase chain reaction cycles comprising a denaturation treatment, wherein hybridized nucleic acid sequences are separated, a hybridization treatment, wherein the target-specific portions of the primary oligonucleotide primers hybridize to the target nucleotide sequences, and an extension treatment,
- 25 wherein the hybridized primary oligonucleotide primers are extended to form primary extension products complementary to the target nucleotide sequence to which the primary oligonucleotide primer is hybridized;
- providing a secondary oligonucleotide primer set characterized by
- (a) a first oligonucleotide primer, having a target-specific portion and a 5'
- 30 upstream secondary primer-specific portion, and (b) a second oligonucleotide primer, having a target-specific portion and a 5' upstream secondary primer-

5

10

blending the primary extension products, the secondary

subjecting the secondary polymerase chain reaction mixture to two

15

20

blending the secondary extension product and the restriction

subjecting the endonuclease digestion reaction mixture to an

25

30

A1

portion of a second oligonucleotide primer of the secondary oligonucleotide primer set, wherein the set of tertiary oligonucleotide primers may be used to amplify all of the secondary extension products;

blending the secondary extension products, the tertiary  
5 oligonucleotide primer set, and the polymerase to form a tertiary polymerase chain  
reaction mixture;

subjecting the tertiary polymerase chain reaction mixture to two or more polymerase chain reaction cycles comprising a denaturation treatment, wherein hybridized nucleic acid sequences are separated, a hybridization treatment, wherein the tertiary oligonucleotide primers hybridize to the secondary extension products, an extension treatment, wherein the hybridized tertiary oligonucleotide primers are extended to form tertiary extension products complementary to the secondary extension products;

providing a plurality of oligonucleotide probe sets, each set  
15 characterized by (a) a first oligonucleotide probe, having a tertiary extension  
product-specific portion and a detectable reporter label, and (b) a second  
oligonucleotide probe, having a tertiary extension product-specific portion,  
wherein the oligonucleotide probes in a particular set are suitable for ligation  
together when hybridized adjacent to one another on a complementary tertiary  
20 extension product-specific portion, but have a mismatch which interferes with  
such ligation when hybridized to any other nucleotide sequence present in the  
sample;

providing a ligase;

blending the tertiary extension product, the plurality of  
oligonucleotide probe sets, and the ligase to form a ligase detection reaction  
mixture;

subjecting the ligase detection reaction mixture to one or more  
ligase detection reaction cycles comprising a denaturation treatment, wherein any  
hybridized oligonucleotides are separated from the tertiary extension products,  
and a hybridization treatment, wherein the oligonucleotide probe sets hybridize at  
adjacent positions in a base-specific manner to their respective tertiary extension  
products, if present, and ligate to one another to form a ligation product sequence  
containing (a) the detectable reporter label and (b) the tertiary extension product-

5

10

15

20

25

30

contacting the ligase detection reaction mixture, after said  
subjecting it to one or more ligase detection reaction cycles, with the solid support  
under conditions effective to hybridize the ligation product sequences to the  
capture oligonucleotides in a base-specific manner, thereby capturing the  
addressable array-specific portions to the solid support at the site with the  
complementary capture oligonucleotide, wherein said detecting indicates the  
presence of ligation product sequences captured using the addressable array-  
specific portions and immobilized to the solid support at particular sites, thereby  
indicating the presence of one or more target nucleotide sequences in the sample.

4. A method according to claim 1, wherein the relative amounts of one or more of a low abundance sequence, differing by one or more single-base changes, insertions, deletions, or translocations and present in a sample in unknown amounts with one or more high abundance sequence in a plurality of target nucleotides, are quantified, said method further comprising:

quantifying, after said subjecting the primary polymerase chain reaction mixture to one or more polymerase chain reaction cycles, the amounts of primary extension products;

providing a known amount of one or more marker target nucleotide sequences;

providing one or more sequence-specific probe sets, including probe sets specifically designed for the marker target nucleotide sequences;

blending the marker target nucleotide sequences, and the probe sets specifically designed for the marker target nucleotide sequences with ligase detection reaction mixture;

quantifying the amount of ligation product sequences; and

comparing the amount of ligation product sequences generated from the unknown low abundance sample with the amount of ligation product sequences generated from known amounts of marker target nucleotide sequences to provide a quantitative measure of the relative level of one or more low abundance target nucleotide sequences in the sample.

5. A method according to claim 4, wherein the relative amounts of one or more of a low abundance sequence, differing by one or more single-base changes, insertions, deletions, or translocations and present in a sample in unknown amounts is present in less than a 1:1,000 molar ratio to the amount of the high abundance sequence in a plurality of target nucleotides.

6. A method according to claim 4, wherein the relative amounts of one or more of a low abundance sequence, differing by one or more single-base changes, insertions, deletions, or translocations and present in a

Variable	Mean	SD	Min	Max	Median	Mode	Skewness	Kurtosis	Shapiro-Wilk	Normality
Age	35.2	12.5	18	65	32	30	0.15	2.8	0.98	Normal
Gender	1.2	0.4	1	2	1	1	0.05	1.2	0.99	Normal
Marital Status	1.5	0.5	1	3	1	1	0.10	1.5	0.97	Normal
Education	12.5	2.5	8	16	12	12	0.20	3.2	0.95	Normal
Income	1500	500	500	3000	1200	1000	0.30	4.5	0.92	Normal
Occupation	1.8	0.6	1	3	1	1	0.12	1.8	0.96	Normal
Health Status	1.2	0.4	1	2	1	1	0.05	1.2	0.99	Normal
Stress Level	2.5	1.0	1	4	2	2	0.25	3.5	0.94	Normal
Life Satisfaction	3.5	1.2	1	5	3	3	0.18	3.0	0.96	Normal
Resilience	2.8	0.8	1	4	2	2	0.22	3.3	0.95	Normal
Optimism	3.2	1.0	1	5	3	3	0.15	2.8	0.97	Normal
Emotional Stability	2.5	0.7	1	4	2	2	0.10	2.5	0.98	Normal
Self-Esteem	3.0	0.9	1	5	3	3	0.12	2.9	0.96	Normal
Life Satisfaction	3.5	1.2	1	5	3	3	0.18	3.0	0.96	Normal
Resilience	2.8	0.8	1	4	2	2	0.22	3.3	0.95	Normal
Optimism	3.2	1.0	1	5	3	3	0.15	2.8	0.97	Normal
Emotional Stability	2.5	0.7	1	4	2	2	0.10	2.5	0.98	Normal
Self-Esteem	3.0	0.9	1	5	3	3	0.12	2.9	0.96	Normal

wherein hybridized nucleic acid sequences are separated, a hybridization treatment, wherein the secondary oligonucleotide primers hybridize to the primary extension products, an extension treatment, wherein the hybridized secondary oligonucleotide primers are extended to form pre-secondary extension products complementary to the primary extension products, wherein the pre-secondary extension products contain one or more nucleotide analogues and opposite strand base changes which facilitate conversion of the primary polymerase chain reaction product sequence into a restriction endonuclease recognition site in the subsequent secondary polymerase chain reaction, wherein the pre-secondary extension products are then used in place of the primary extension products in the secondary polymerase chain reaction mixture.

9. A method according to claim 8, wherein the oligonucleotide probes in a particular oligonucleotide probe set have a unique length so that the ligation product sequences which they form can be distinguished from other nucleic acids:

separating the ligation product sequences by electrophoretic mobility prior to said detecting and distinguishing, after said detecting, the ligation product sequences which differ in electrophoretic mobility.

10. A method according to claim 8, wherein the second oligonucleotide probe of each set has an addressable array-specific portion, said method further comprising:

providing a solid support with different capture oligonucleotides immobilized at different particular sites, wherein the capture oligonucleotides have nucleotide sequences complementary to the addressable array-specific portions and

contacting the ligase detection reaction mixture, after said subjecting it to one or more ligase detection reaction cycles, with the solid support under conditions effective to hybridize the ligation product sequences to the capture oligonucleotides in a base-specific manner, thereby capturing the addressable array-specific portions to the solid support at the site with the

5

15

30

[illegible]

13. A method according to claim 11, wherein the relative amounts of one or more of a low abundance sequence, differing by one or more single-base changes, insertions, deletions, or translocations and present in a sample in unknown amounts is present in a molar ratio of less than 1:10,000 to the amount of the high abundance sequence in the sample.

14. A method according to claim 11, wherein the relative amounts of one or more of a low abundance sequence, differing by one or more single-base changes, insertions, deletions, or translocations and present in a sample in unknown amounts is present in a ratio molar ratio of less than 1:100,000 to the amount of the high abundance sequence in the sample.

15. A method according to claim 8, where the nucleotide analog of at least one oligonucleotide primer of the pre-secondary oligonucleotide primer set is at the 3' end of the primer.

16. A method according to claim 8, where the nucleotide analog is selected from the group consisting of 1-(2'-deoxy-β-D-ribofuranosyl)imidazole-4-carboxamide, 1-(2'-deoxy-β-D-ribofuranosyl)-3-nitropyrrole, 2'-deoxyinosine, 6-(2'-deoxy-β-D-ribofuranosyl)-6H,8H-3,4-dihydropyrimido[4,5-c][1,2]oxazine-7-one, 2-amino-7-(2'-deoxy-β-D-ribofuranosyl)-6-methoxyaminopurine, 1-(2'-deoxy-β-D-ribofuranosyl)-4-iodopyrazole, 1-(2'-deoxy-β-D-ribofuranosyl)pyrrole-3-carboxamide, and 1-(2'-deoxy-β-D-ribofuranosyl)-4-nitropyrrole.

17. A method according to claim 1 further comprising:  
repeating the endonuclease digestion reaction after said subjecting the tertiary polymerase chain reaction mixture to two or more polymerase chain reaction cycles and after said subjecting the ligase detection reaction mixture to one or more ligase detection reaction cycles, wherein, during said repeating the endonuclease digestion reaction, the restriction endonuclease recognizes and cleaves the restriction endonuclease recognition site contained within any

a2  
~~remaining-high abundance target, thereby selectively destroying the high abundance tertiary extension products.~~

18. A kit for identifying one or more low abundance sequences
- 5 differing by one or more single-base changes, insertions, or deletions, from a high abundance sequence in a plurality of target nucleotide sequences comprising:
- providing a primary oligonucleotide primer set characterized by (a)
- a first oligonucleotide primer containing a target-specific portion, and (b) a second
- 10 oligonucleotide primer containing a target-specific portion, wherein the primary oligonucleotide primers are suitable for hybridization on complementary strands of a corresponding high and low abundance target nucleotide sequences to permit formation of a primary extension product, but have a mismatch which interferes with formation of such a polymerase chain reaction product when hybridized to any other nucleotide sequence present in the sample;
- 15 providing a secondary oligonucleotide primer set characterized by (a) a first oligonucleotide primer, having a target-specific portion and a 5' upstream secondary primer-specific portion, and (b) a second oligonucleotide primer, having a target-specific portion and a 5' upstream secondary primer-specific portion, wherein the secondary oligonucleotide primers in a particular set
- 20 are suitable for hybridization on complementary strands of the primary extension products to permit formation of a secondary extension product which contains or creates a restriction endonuclease recognition site when amplifying the high abundance target, but does not contain or create a restriction endonuclease recognition site when amplifying the one or more low abundance targets;
- 25 providing a tertiary oligonucleotide primer set characterized by (a) a first tertiary primer containing the same sequence as the 5' upstream portion of the first oligonucleotide primer of the secondary oligonucleotide primer set, and (b) a second tertiary primer containing the same sequence as the 5' upstream portion of a second oligonucleotide primer of the secondary oligonucleotide
- 30 primer set, wherein the set of tertiary oligonucleotide primers may be used to amplify all of the secondary extension products; and
- providing a plurality of oligonucleotide probe sets, each set characterized by (a) a first oligonucleotide probe, having a tertiary extension

product-specific portion and a detectable reporter label, and (b) a second oligonucleotide probe, having a tertiary extension product-specific portion, wherein the oligonucleotide probes in a particular set are suitable for ligation together when hybridized adjacent to one another on a complementary tertiary extension product-specific portion, but have a mismatch which interferes with such ligation when hybridized to any other nucleotide sequence present in the sample.

19. A kit according to claim 18 further comprising:  
providing a pre-secondary oligonucleotide primer set characterized by (a) a first oligonucleotide primer, having a target-specific portion, and (b) a second oligonucleotide primer, having a target-specific portion, wherein the target-specific portions are identical or substantially identical to the secondary oligonucleotide primer set but at least one primer contains one or more nucleotide analogs, wherein the pre-secondary oligonucleotide primers in a particular set are suitable for hybridization on complementary strands of the primary extension products to permit formation of a pre-secondary extension product which contains one or more nucleotide analogs and opposite strand base changes, wherein the pre-secondary oligonucleotide primer set facilitates conversion of the primary extension product sequence into a restriction endonuclease recognition site in a subsequent secondary polymerase chain reaction.

20. A kit according to claim 18 further comprising:  
a ligase.

21. A kit according to claim 18 further comprising:  
a polymerase.

22. A kit according to claim 18 further comprising:  
a restriction endonuclease.